

Abundances of metal-weak thick-disc candidates

P. Bonifacio, M. Centurion and P. Molaro

Osservatorio Astronomico di Trieste, Via G.B. Tiepolo 11 34131, Trieste – Italy

ABSTRACT

High resolution spectra of 5 candidate metal-weak thick-disc stars suggested by Beers & Sommer-Larsen (1995) are analyzed to determine their chemical abundances. The low abundance of all the objects has been confirmed with metallicity reaching $[\text{Fe}/\text{H}] = -2.9$. However, for three objects, the astrometric data from the Hipparcos catalogue suggests they are true halo members. The remaining two, for which proper-motion data are not available, may have disc-like kinematics. It is therefore clear that it is useful to address properties of putative metal-weak thick-disc stars only if they possess full kinematic data. For CS 22894-19 the abundance pattern similar to those of typical halo stars is found, suggesting that chemical composition is not a useful discriminant between thick-disc and halo stars. CS 29529-12 is found to be C enhanced with $[\text{C}/\text{Fe}] = +1.0$; other chemical peculiarities involve the *s* process elements: $[\text{Sr}/\text{Fe}] = -0.65$ and $[\text{Ba}/\text{Fe}] = +0.62$, leading to a high $[\text{Ba}/\text{Sr}]$ considerably larger than what is found in more metal-rich carbon-rich stars, but similar to LP 706-7 and LP 625-44 discussed by Norris et al. (1997a). Hipparcos data have been used to calculate the space velocities of 25 candidate metal-weak thick-disc stars, thus allowing us to identify 3 *bona fide* members, which support the existence of a metal-poor tail of the thick-disc, at variance with a claim to the contrary by Ryan & Lambert (1995).

Key words: Stars: abundances – Stars: Population II – Stars: fundamental parameters – Galaxy: halo

1 INTRODUCTION

The disc of our Galaxy is rotationally supported with a small velocity dispersion and a scale height of about 300 pc. The existence of a “thick disc” was highlighted by Gilmore & Reid (1983) using star count data. The proposed scale heights are in the range 1.0-1.5 kpc, the asymmetric drift is in the range of 30-50 kms^{-1} , and the velocity dispersion is about twice that of the thin disc. Discs with a vertical scale height in excess of 1 kpc may be seen in external edge-on spiral galaxies with small bulges, similar to our own (Van der Kruit & Searle 1981a, 1981b, 1982). Because suggested kinematical and chemical parameters of the thick disc are similar to those of the thin disc, this makes their determination prone to selection effects, and the issue of discreteness difficult to establish or reject. Whether the thick disc is a discrete component of the Galaxy or it is merely an “extended” configuration of the disc, as suggested by Norris & Ryan (1991), is still an open question. For the sake of clarity, in the following we shall refer to the two as “thin” (scale height $\approx 300\text{pc}$) and “thick” (scale height $>1\text{ kpc}$) without implying by this the discreteness of the components. The origin of the thick disc is not clear. In their review Gilmore, Wyse & Kuijken (1989) list the possible formation mechanisms, which fall into three broad categories: 1) slow formation of the disc ; 2) rapid formation of the thin disc and successive heating ; 3) accretion of satellite galaxies.

The thick disc is probably as old as the halo: Wyse & Gilmore (1995) claim that the thick disc is “apparently exclusively old”. The absence of RR Lyrae variables with disc kinematics would rule out that the disc is as old as the globular clusters, as had been noted by Majewski (1992). However the recent work of Martin & Morrison (1998) using new kinematic data for a sample of 130 nearby RR Lyrae variables, was able to isolate a subsample of 21 stars with $[\text{Fe}/\text{H}] \leq -1.0$ and disc-like kinematics. The work of Fuhrmann (1998a), which makes use of Hipparcos data and high resolution, high S/N spectra, of about 50 nearby F and G type stars, supports the notion that the thick disc is as old as the halo ($\approx 12\text{Gyr}$), while the thin disc is younger than 9-10 Gyr, thus highlighting a hiatus in star-formation in the Galaxy.

The existence of the thick disc and its kinematical parameters have a direct bearing on the interpretation of data aimed at the derivation of the rotational properties of the halo. As was pointed out by Majewski (1992) “halo tracer surveys select stars from regions dominated by either the thin or thick disc”. A mixture of rapidly rotating disc stars and retrograde rotating halo stars could explain why such surveys determine a low prograde rotation for the halo, while Majewski’s high *z* sample reveals a sizeable retrograde motion.

The kinematical distinction between disc (thin or thick) and halo is rather abrupt and should not give rise to any confusion. However, if we consider a metallicity biased sample of faint stars, their kinematics are imperfectly known (in general only radial velocity and photometric or spectroscopic parallax are available), making it impossible to assign unambiguous membership to the disc or to the halo on kinematical grounds. Can a selection may be made on chemical grounds alone? An answer to this question requires knowledge of the metallicity distribution of the thick disc. The metallicity of the thick disc is considered to be “intermediate” between thin disc and halo (Gilmore & Wyse, 1985; Carney Latham & Laird 1989) but a well defined metallicity distribution is not yet available. Given the difficulty of isolating a pure thick disc sample, it is not surprising that different selection criteria lead to different answers. Several studies have found a mean metallicity of the thick disc between -0.6 and -0.4 (Gilmore & Wyse 1985, Sandage & Fouts 1987; Yoss, Neese & Hartkopf 1987, Carney et al. 1989; Friel 1987, Hartkopf & Yoss 1982; Ratnatunga & Freeman 1989). Morrison, Flynn & Freeman (1990) claimed that a metal-weak tail of the thick disc extends down to $[\text{Fe}/\text{H}] = -1.6$. On the other hand, Majewski (1992) finds a mean metallicity of -0.13 , but suggests that the distribution is markedly non-gaussian and has a metal weak tail even more extreme than suggested by Morrison et al. (1990). Gilmore & Wyse (1995) find from their *in situ* sample that the metallicity distribution peaks at $[\text{Fe}/\text{H}] = -0.7$ and there is a large overlap in metallicity between thin and thick disc, the thin disc extending down to at least -0.75 and the thick disc up to at least -0.3 . Reid et al. (1997), studied a sample of faint M dwarfs at high galactic latitude and found their data to be consistent with star-count models including three components (thin disc, thick disc, halo) and that up to 2 kpc above the plane the abundances are close to the values of the old thin disc. The analysis of Reid (1998), who used a recalibration of the colour magnitude diagrams of globular clusters based on local subdwarf calibrators with Hipparcos data, to define a reference grid in the $(M_V, B - V)$ plane, implies that $\approx 15\%$ of local disc stars have $[\text{Fe}/\text{H}]$ below -0.3 . Beers & Sommer-Larsen (1995) suggest that stars with disc-like kinematics may be found for arbitrarily low abundances, although their sample, selected mainly on the criterion of low metal abundance, is not well suited to determine the metallicity distribution of the thick disc. In their paper Beers & Sommer-Larsen provided a list of likely metal-poor thick disc stars, i.e. stars whose position is such that the line of sight component of the thick disc rotational velocity is in excess of 100 km s^{-1} .

Given the probable similarity in age of the halo and thick-disc, the existence of a metal-weak tail should be expected. However the existence of a metal-weak tail of the thick disc is seriously challenged by the results of Ryan & Lambert (1995), who analyzed at high spectral resolution a sample of expected metal-poor disc stars taken from the samples of Norris et al. (1985) and Morrison et al. (1990) and found no star with disc kinematics below $[\text{Fe}/\text{H}] = -1.0$, arguing that the metallicities inferred from DDO photometry by Morrison et al. (1990) were underestimated.

Nissen & Schuster (1997) studied a sample of stars of intermediate metallicity belonging to both the disc and halo for which their lowest metallicity disc star has $[\text{Fe}/\text{H}] = -1.3$. In the Fuhrmann (1998a) sample the most metal weak thick disc star is around $[\text{Fe}/\text{H}] = -1.0$. He was able to show that a discrimination between thin and thick disc is possible, if a combination of kinematics, abundance and age data is used.

Here we report on the detailed abundance analyses of 5 of the most interesting candidates, based on high resolution spectra. Li abundances for two of these objects have already been reported by Molaro, Bonifacio & Pasquini (1997).

2 OBSERVATIONS AND DATA REDUCTION

In September 1996 we observed CS 29529-012, CS 22894-019, BD $-14^\circ 5890$, HD165195, HD26169 and CS 22959-007 with the CASPEC spectrograph at the ESO 3.6 m telescope. The width of the lines of CS 22959-007 revealed that it is a spectroscopic binary, it has been therefore dropped from the present analysis and we plan to collect other observations to elucidate its nature. The stars which were selected from Table 8 of Beers & Sommer-Larsen, had already been observed by us in the red (Molaro Bonifacio & Pasquini 1997). The echelle grating with $31.6 \text{ lines mm}^{-1}$ and the Long Camera (f/3) were used. The detector was the Tek CCD 1024×1024 with 24 micron wide pixels and a readout noise of about $4e^-$ per pixel. With the Long Camera the scale at the detector was about $0.059 \text{ \AA /pixel}$ at 473.6 nm . The slit was set at $1.2''$ arc seconds providing a resolving power of $R = \lambda/\delta\lambda \approx 30000$, as measured from the Thorium-Argon emission lines. The echellograms were reduced using the MIDAS ECHELLE package. The background was subtracted and cosmic rays removed with the FILTER/ECHELLE command. The images were not flat-fielded since this process increases the noise in the spectra and, at the wavelengths covered by our data, fringing is not important. The spectra were then wavelength calibrated using the Th-Ar lamp spectrum, the root mean square error of the calibration was of the order of $0.2 - 0.3 \text{ pm}$ for all the 36 orders observed on the detector. The spectrum of CS 29529-012 and one of the spectra of CS 22894-019 were taken in bright time and the sky spectrum was subtracted. The other spectrum of CS 22894-019 was taken during a total moon eclipse and sky subtraction was not necessary. The spectra of the other significantly brighter stars were not seriously affected by the sky, because the exposure times were much shorter. For CS 29529-012 and CS 22894-019 we also made use of the red spectra described in Molaro, Bonifacio & Pasquini (1997).

A synthetic spectrum was computed for CS 29529-012 and CS 22894-019 using the atmospheric parameters derived by Molaro et al. (1997) and over-plotted on the observed spectrum for identification purposes. Blended or noisy lines were removed in order to define a preliminary list of *bona fide* lines to be measured. These lines were measured on all the available spectra by fitting a gaussian with the IRAF task `splot` which allowed the determination of line positions and equivalent

Table 1. Basic data of program stars

Star	T_{eff}	log g spec.	log g Hip.	[Fe/H]	[M/H] ^a	ξ kms ⁻¹	v_r kms ⁻¹
CS 22894-19	6060	3.5	-	-2.90	-3.00	2.0	32 ± 2
CS 29529-12	6197	3.0	-	-2.27	-2.00	1.7	93 ± 2
BD -14°5890	4700 ^b	1.4	2.34	-2.52	-2.50	1.9	118 ± 2
HD 165195	4500 ^c	1.1	1.42	-1.92	-2.00	1.6	0 ± 2
HD 26169	5000 ^d	2.0	2.46	-2.61	-2.50	1.9	20 ± 15

^a metallicity of the adopted model

^b Cavallo et al (1997)

^c Francois (1996)

^d Peterson et al (1990)

widths. Since the dark-time spectrum of CS 22894-019 was by far superior to the bright-time one, we used only the equivalent widths measured from this one. However the line positions were taken from both and the resulting radial velocities were averaged.

3 ATMOSPHERIC PARAMETERS

For the stars BD -14°5890, HD165195 and HD26169 we adopted effective temperatures from the literature, as detailed in Table 1, while for CS 29529-012 and CS 22894-019 we adopted the temperature derived from $(B-V)_0$ by Molaro et al. (1997).

For all the stars the gravity was derived from the FeI/FeII ionization equilibrium and the microturbulent velocity by the constraint that the FeI abundance be independent of equivalent width. Given the poor S/N of the spectra of the CS stars we were forced to use mainly strong lines and therefore the microturbulent velocity is poorly constrained.

Both CS 22894-19 and CS 29529-12 are classified as dwarfs by Beers et al. (1992). For CS22894-19 we derived a gravity of $\log g = 3.5$ using 3 Fe II lines. We had excluded the FeII 516.9 nm line because it yields an abundance which is almost 0.5 dex lower than that of the other three lines, regardless of the gravity. By taking the Fe II abundance to be the average of all four lines one would derive a gravity of $\log g = 4.0$. Since we assumed this line to be affected by noise we removed it from the analysis. Further support for the subgiant status of CS22894-19 comes from the Mg I b triplet lines: with $\log g = 3.5$ the computed lines provide a satisfactory fit, whereas with a $\log g = 4.0$ the computed lines are considerably broader than the observed ones.

Our gravity for CS29529-12 was $\log g = 3.0$ derived using five Fe II lines. Both Fe I and Fe II have a rather small rms scatter of about 0.2 dex in the abundances. Since a change as large as 0.5 dex in $\log g$ does not produce a difference larger than 0.2 dex in the abundances of the two species, we consider the gravity to be uncertain by ± 0.5 dex. With the $\log g = 3.00$ the computed line strengths of the b triplet are internally consistent and match the observed one. However the Mg I 470.2991 nm line yields an abundance which is about 0.5 dex larger. A decrease of surface gravity or microturbulent velocity lessens the disagreement among Mg lines. For $\log g = 2.5$ and $\xi = 1.5$, values which are not unreasonable, the difference is 0.37 dex, and for $\log g = 2.0$ $\xi = 1.5$ it is 0.27 dex. This is because the Mg b triplet lines are sensitive both to pressure and to microturbulence, while the abundance from the Mg I 470.2991 nm line is not very sensitive to gravity, where a decrease of 1 dex in $\log g$ produces only an increase of 0.065 dex in Mg abundance.

Recently it has been realized that ionization equilibria provide gravities which are up to 0.5 dex lower than the gravities derived in stars with accurate Hipparcos parallaxes (Nissen Hoeg & Schuster 1997; Fuhrmann 1998b). We preferred the $\log g$ given by ionization equilibrium for abundance analysis, both for homogeneity with the analysis of the two stars for which Hipparcos data are not available, and because the structure of the atmospheric model satisfying the iron ionization equilibrium better reproduces the observed spectrum than does a model with the Hipparcos gravity.

3.1 Evolutionary status

To gain some insight into the evolutionary status of the stars in our sample we compared their temperatures and gravities with those predicted by theoretical isochrones. In Figure 1 we show the $T_{\text{eff}} - \log g$ diagram with the Revised Yale Isochrones (RYI ; Green et al 1987) for metallicity $[\text{Fe}/\text{H}] = -2.0$ and -3.0 , an age of 12 Gyr and helium mass fraction $Y=0.25$. The isochrones of the Padova group (Bertelli et al 1994) are also shown for $[\text{Fe}/\text{H}] = -1.7$ ($z=0.0004$), ages of 12 Gyr and 6 Gyr, and $Y=0.23$.

The position of the two CS stars is anomalous, although given the large uncertainty associated with their gravities they are in fact compatible with a TO or SG status. They actually occupy an area in the diagram occupied by Horizontal Branch




Figure 1. T_{eff} -log g diagram for our program stars. Solid lines are RYI isochrones for $[\text{Fe}/\text{H}] = -2.0$ and -3.0 (bluest), 12 Gyr and $Y=0.25$. The dots represent the Padova isochrones for $[\text{Fe}/\text{H}] = -1.7$, 12 Gyr and 6 Gyr (bluest) and $Y=0.23$. The age-metallicity degeneracy is evident.

stars, or Subgiant stars of much younger age (below 8 Gyr). For the three giant stars, for which Hipparcos parallaxes are available, we may compute gravities using the equation

$$\log \frac{g}{g_{\odot}} = \log \frac{M}{M_{\odot}} + 0.4(V_0 + B.C.) + 2 \log \pi + 0.12$$

(see e.g. Nissen et al 1997), where we assumed $M = 0.8M_{\odot}$, as is appropriate for giant stars on the RYI isochrone for $[\text{Fe}/\text{H}] = -2.0$, shown in figure 1 (note that changing the mass to $0.5M_{\odot}$ would decrease the $\log g$ by only 0.2 dex). These gravities are, as expected, larger than the spectroscopic gravities and are given in table 1. The HB hypothesis may be discarded, because both stars show a normal Li abundance (Molaro et al. 1997) while a star during its red giant phase is expected to decrease its photospheric Li content due to depletion and dilution. That the stars are very young is rather problematic considering their low abundance. Moreover the results of Fuhrmann (1998a) imply that the thick disk is an old population. From our spectroscopic gravities we may derive distances for these stars, which are much greater than those given by Beers et al. (1992), who assumed a TO status. For CS22894-19 we derive 1.7 kpc, compared to 1.1 kpc of Beers et al.; for CS29529-12 1.2 kpc, compared to 0.4 kpc of Beers et al.

It is for CS 29529-12 that the situation is more critical: if the ionization equilibrium gravity is wrong then the star can really be a turn-off star. If this is the case, then the surface gravity is around $\log g = 4.0$, whichever isochrone set we use. However, on performing an abundance analysis with this surface gravity we find several chemical inconsistencies, as shall be detailed in section 5. There will be a discrepancy of 0.34 dex between the results for Fe I and Fe II, which is not inconsistent with the rms scatter of each species. The chemical composition would then show some peculiar ratios: $[\text{Mg}/\text{Fe}] = +0.17 \pm 0.27$, $[\text{Ca}/\text{Fe}] = +0.13 \pm 0.26$ and $[\text{TiII}/\text{FeII}] = +0.40 \pm 0.29$. The errors are estimated by just summing under quadrature the line to line scatter given in Table 3. The errors in T_{eff} and ξ for $[\text{Mg}/\text{H}]$, $[\text{Ca}/\text{H}]$, $[\text{Ti}/\text{H}]$ and $[\text{Fe}/\text{H}]$ are in the same direction and about the same magnitude, so that they will cancel out. The errors in $\log g$, instead, are in different directions for $[\text{Mg}/\text{H}]$, $[\text{Ca}/\text{H}]$ and $[\text{Fe}/\text{H}]$, so that this contribution should also be added under quadrature, making the errors on $[\text{Mg}/\text{Fe}]$, and $[\text{Ca}/\text{Fe}]$ even larger. For Ti we considered only the $[\text{TiII}/\text{FeII}]$ so that errors in $\log g$ will cancel out. Although the errors allow all these ratios to be consistent with a “standard” $[\alpha/\text{Fe}] = +0.4$, their values, taken at face value, appear anomalous. While the existence of α -poor metal-poor stars has been reported by Carney et al (1997), Nissen & Schuster (1997) and Jehin et al (1997), it is difficult to believe that Ti is over-enhanced with respect Mg and Ca. In fact while Mg and Ca are “pure” α -elements, Ti is also produced in nuclear statistical equilibrium together with iron-peak elements, thus it is expected to be under-enhanced with respect to Mg and Ca (Timmer et al. 1995). Observers have either claimed that Ti goes in lockstep with Ca while Mg is over-enhanced (Primas et al. 1994, Ryan et al. 1996), or that Ti is slightly under-enhanced with respect to Ca and Mg (McWilliam et al. 1995). So far there has been no report of a Ti over-enhancement in metal-poor stars.

4 MODEL ATMOSPHERES AND ANALYSIS

For each star we computed a model-atmosphere using version 9 of the ATLAS code (Kurucz 1993) with the above determined parameters. We used the α -enhanced opacity distribution functions (ODFs), a microturbulence velocity of 1 km s^{-1} and no convective overshooting. These model atmospheres and the measured equivalent widths were used as input to the version 9 of the WIDTH code (Kurucz 1993) to determine the elemental abundances. At this stage some of the lines were removed from the analysis when their abundance was deviant from that of the other lines of the same species by more than 1σ . Equivalent widths, atomic data and derived abundances for the lines which were retained are listed in Table 2.

We consider errors which arise from errors in the equivalent widths and errors in the model-atmosphere parameters. We ignore errors in the atomic data and errors due to shortcomings in the model-atmosphere (treatment of convection, reliability of 1D-models). Formula (7) of Cayrel (1988) provides a conservative estimate of the errors in the equivalent widths: for our instrumental setup, we estimate an error of 0.5 pm for the dark-time spectrum of CS22894-019, 0.8 pm for the spectrum of CS29529-012 and around 0.1 pm for the giants. Misplacement of the continuum can introduce an error in the equivalent width of up to 1.6 pm. Somewhat smaller errors are expected for the lines which could be measured in two adjacent orders. We believe that a realistic estimate of abundance errors may be obtained from the line to line scatter when several lines are available. We investigate in detail the errors only for CS29529-012. For each element in Table 3 we report five errors: σ is the r.m.s. error or line to line scatter for species with more than one line, the other columns are the variation in abundance deduced for an increase in $\log g$ by 0.5 dex, a decrease in T_{eff} by 130 K, an increase in ξ by 0.3 km s^{-1} and an increase in equivalent widths by 0.4 pm. These errors may be used as a guide to assess the influence on the abundances of errors in model parameters and equivalent widths. We expect the situation for the other stars to be quite similar.

5 RESULTS

For CS 22894-19 and CS 29529-12 we derived $[\text{Fe}/\text{H}] = -2.90$ and -2.27 respectively, the metallicities derived from the Ca II K index by Beers et al. (1992) for the two stars are -3.03 and -1.54 , respectively. While for CS 22894-19 there is a substantial agreement, for CS 29529-12 our metallicity is considerably lower than previously estimated. A stronger than normal Ca II K line, or contamination of an interstellar Ca II line, could have explained the discrepancy between our metallicity and that of Beers et al. (1992). However we verified with spectrum synthesis that the Ca abundance derived from the Ca I lines is compatible with the Ca II H and K lines.

For a few lines in the giant stars we resorted to spectrum synthesis either because the lines are mildly blended or because we felt that the gaussian fit used to measure the equivalent width did not capture satisfactorily the line wings as is the case for the Mg I b triplet lines of HD 161695.

For Mn we report the abundances only for the two CS stars, because for the giants the lines are saturated, and thus not suitable for abundance determination. The lines were treated by spectrum synthesis accounting for the hyperfine splitting. The relevant atomic data was taken from McWilliam et al. (1995).

The abundances in CS 22894-19 are similar to those of typical halo stars of comparable metallicity and in that respect unremarkable.

CS29529-12 deserves some additional remarks. Among the observed Cr lines there is some discrepancy. The abundance of Cr given in Table 4, has been derived from one resonance line of multiplet 1, 427.5 nm (we excluded CrI 425.4 nm because it is rather large in our spectrum and we suspect it to be blended, possibly with a Cr II line, and 429.0 nm because it yields an abundance which is 1 dex higher than the other lines) and the three lines of multiplet 7. All the $\log gf$ values are computed from the transition probabilities of the compilation of Younger et al (1978). The data for multiplet 7 are less accurate, owing to a possible self-absorption in the emission-line experiments; however their uncertainties should be within 50%, which translates into a 0.2 dex uncertainty in $\log gf$ and therefore abundance.

The Sr abundance is derived from the Sr II 421.5 nm resonance line only, because the stronger Sr II 407.8 nm line is affected by noise in our spectrum. The barium abundance is obtained from the 445.5 nm line. This line may be potentially affected by hyperfine structure (*hfs*) but in an amount which depends on the relative importance of *s* or *r* processes (see Norris et al 1997a). When the *s* process is dominant there is a minimal production of the odd isotopes where the *hfs* is important.

Sr is over-deficient by -0.65 dex, while Ba is over-abundant by 0.62 dex. This yields $[\text{Ba}/\text{Sr}] = +1.3$, which is rather unusual for “normal” metal-poor stars, where it is found negative. However *s*-enriched metal-poor stars, such as carbon-rich stars (CH,sgCH), do show $[\text{Ba}/\text{Sr}]$ ratios in the range 0.5 to 2, this is generally understood to be due to the fact that at low metallicities the main neutron source necessary for the *s*-process is $^{13}\text{C}(\alpha, n)^{16}\text{O}$, rather than $^{22}\text{Ne}(\alpha, n)^{25}\text{Mg}$ (Luck & Bond 1982). In fact CS 29529-12 is mildly C enhanced with $[\text{C}/\text{Fe}]$ on the order of $+1.0$. The carbon abundance has been derived from spectral synthesis of the G band. The star has not been noticed to have extremely strong G-band by Beers et al. (1992), as its GP index is 0.92, which is only slightly larger than the median GP value for metal-poor stars around metallicity -2.0 .

We do not have information on nitrogen and oxygen since the CN 388.3 nm band falls outside our spectral range and the noise at the OI 630.0 nm line does not allow us to place useful limits on the oxygen abundance. Adopting $[\text{O}/\text{Fe}] = +0.6$ dex

Table 2. line data and individual abundances

	$\lambda(\text{nm})$	$\chi(\text{cm}^{-1})$	loggf	$EW(\text{pm})$ CS 22894-19	ϵ	$EW(\text{pm})$ CS 29529-12	ϵ	$EW(\text{pm})$ BD $-14^\circ 5890$	ϵ	$EW(\text{pm})$ HD 165195	ϵ	$EW(\text{pm})$ HD 26169	ϵ
Na I	588.9951	0.000	0.117	6.57	3.15	19.10	5.57						
Na I	589.5924	0.000	-0.184	4.47	3.13	17.20	5.68						
Mg I	470.2991	35051.264	-0.666	2.61	5.33	7.36	6.21	9.72	5.92	—	-6.20	8.08	5.76
Mg I	517.2684	21870.464	-0.402	11.26	4.87	14.50	5.71	27.00	5.51	—	-6.40	21.12	5.31
Mg I	518.3604	21911.178	-0.180	13.20	4.92	15.60	5.63	30.56	5.43	—	-6.40	24.52	5.31
Al I	396.1520	112.061	-0.323	6.98	3.01	9.28	3.70	28.25	3.14	22.82	3.90	13.62	3.16
Ca I	422.6728	0.000	0.243	10.75	3.49	12.63	4.26	30.02	4.15	—	—	23.13	4.12
Ca I	430.2528	38551.558	0.275	4.48	4.01	5.69	4.37	—	—	—	—	—	—
Ca I	445.4779	37757.449	0.252	3.84	3.91	5.50	4.35	8.98	4.05	—	—	7.96	4.01
Ca I	616.2173	31539.495	0.100	—	—	6.52	4.65	—	—	—	—	—	—
Sc II	424.6822	2540.950	0.322	3.49	0.07	7.24	0.74	12.51	0.42	15.83	1.12	10.89	0.44
Ti II	430.0049	9518.060	-0.750	3.75	2.39	—	—	—	—	—	—	—	—
Ti II	439.5033	8744.250	-0.650	5.00	2.41	9.30	3.31	—	—	21.34	3.89	10.17	2.47
Ti II	444.3794	8710.440	-0.810	—	—	6.80	2.87	11.22	2.53	14.59	3.33	10.18	2.63
Ti II	446.8507	9118.260	-0.600	3.90	2.21	7.10	2.76	12.65	2.69	16.90	3.48	10.83	2.61
Ti II	450.1273	8997.710	-0.750	—	—	8.30	3.17	12.00	2.69	16.08	3.50	10.07	2.56
Ti II	457.1968	12676.970	-0.530	3.61	2.49	7.00	3.07	11.91	2.93	16.86	3.82	10.17	2.85
Cr I	425.4332	0.000	-0.114	3.99	2.49	—	—	11.66	2.67	16.10	3.35	9.79	2.57
Cr I	427.4796	0.000	-0.232	4.40	2.68	6.55	3.33	12.15	2.89	17.07	3.64	10.48	2.86
Cr I	428.9716	0.000	-0.365	2.18	2.36	9.94	4.35	—	—	—	—	—	—
Cr I	520.4506	7593.160	-0.170	1.81	2.90	3.20	3.41	—	3.17	—	3.50	—	—
Cr I	520.6038	7593.160	0.030	2.05	2.77	5.50	3.66	—	2.78	—	—	—	2.72
Cr I	520.8419	7593.160	0.160	2.13	2.66	4.40	3.31	—	2.98	—	3.50	—	—
Mn I	403.0753*	0.000	—	—	2.20	—	—	—	3.00	—	—	12.04	2.87
Mn I	403.3062*	0.000	—	—	2.15	—	—	—	—	—	—	10.17	2.50
Mn I	403.4483*	0.000	—	—	2.20	—	—	—	3.12	—	—	—	—

* treated by spectrum synthesis with HFS splitting as given by McWilliam et al (1995)

which is typical of Population II stars we have $[C/O]=+0.4$, which is not enough to classify it as a carbon star, but which clearly shows a carbon enhancement with respect to normal dwarfs.

The cases of carbon-enhanced metal-poor stars are discussed by Norris et al. (1997a) who emphasized the relatively high frequency of carbon-enhanced stars at low metallicity compared to current chemical galactic evolution models. These stars show a large variety of chemical behaviour for the neutron capture elements going from the case of CS 22892-52, where a large abundance of r elements is present, to the case of CS 22957-027 (Norris et al 1997b, Bonifacio et al 1998) where the C overabundance is observed together with a strong underabundance in the neutron-capture elements.

The distinctive peculiarity of CS 29529-12 is that the carbon enhancement is observed together with a Ba enhancement, but with very little Sr. The quantity $[hs/ls]$, defined as the difference between the mean abundance of the heavy s process elements, such as Ba, and the abundance of the lighter ones such as Sr, is heavily weighted toward higher atomic number in stars which show neutron capture elements enhancement.

Norris et al. found that the $[hs/ls]$ ratio increases at low metallicities being about 1.5 for the two stars LP 625-44 and LP 706-7 which have $[Fe/H]$ around -3.0 . CS 29529-12 is slightly more metal-rich and with a less extreme carbon abundance but an $[hs/ls]$ ratio very similar to that of these two stars.

Chemical peculiarities in cool stars are interpreted to be the result either of mixing in the star itself or of accretion from an evolved companion. In particular the formation of Ba stars, CH stars and dwarf carbon stars probably involves the evolution of a binary system.

We further note, with regard to chemical peculiarities of CS 29529-12, that the lithium abundance ($\epsilon(\text{Li}) = \log(\text{Li}/\text{H}) + 12 = 2.21$) is found at the value typical of the other low-metallicity turn-off stars (Molaro et al 1997). If the star had undergone some unusual surface disturbance as a result of self-mixing or of the dredge up of processed material onto their surfaces, the Li surface abundance would have been modified. Li is easily destroyed if mixed into deeper layers where temperatures reach a few million degrees. The apparent normality of Li is thus suggestive evidence that the star has not experienced major modifications of its surface abundance by mixing with internal layers. A similar case is that of LP 706-7 (= G268-32) with $\epsilon(\text{Li}) = 2.25$ (Thorburn 1994) and CS 22898-027 with $\epsilon(\text{Li}) = 2.52$ (Thorburn & Beers 1992)

The binarity hypothesis remains favoured and should be checked by additional observations. While, for LP 625-44, radial velocity and Li abundance are consistent with the binarity for LP 707-7 there is no clear evidence of radial velocity variations and its Li abundance is at the level of the *Spite Plateau*. It would be interesting to establish if CS 29529-12 is analogous to LP 706-7, because these objects are very difficult to account for in present stellar evolution theory.




Figure 2. Na I D lines of CS 29529-12.

The abundance of Na deduced from the D lines is remarkably high, the lines are shown in Fig. 2. It is true that in this temperature and metallicity range the sodium D lines are known to be strongly affected by NLTE effects (Baumüller, Butler & Geheren, 1998), however the correction should be at most of the order of -0.5 dex in $[\text{Na}/\text{Fe}]$, thus leaving an uncomfortably large overabundance of 1.0 dex. For confirmation of the overabundance we searched for the subordinate lines at 568.2633 nm, 568.82 nm and 616.0747 nm. The last is affected by atmospheric emission in our spectrum, the first two should have an equivalent width around 4 pm for the Na abundance implied by the resonance lines, one is marginally detected but the other is not, so the test is inconclusive. The sodium overabundance persists even if we adopt the gravity of $\log g = 4.0$, although reduced by 0.6 dex. The three bright giant stars have been studied by several authors, and so in the appendix we compare our results with the literature.

6 KINEMATICS OF METAL WEAK THICK DISC CANDIDATES

For all the program stars we determined radial velocities from the lines used in the abundance analysis. The r.m.s scatter is on the order of 0.7 - 0.9 kms^{-1} for the brighter giant stars, while it is of the order of 1.6 kms^{-1} for the lower S/N data of the fainter CS stars. In both cases the resulting error in the radial velocity is dominated by the uncertainty in the zero point shift, which is on the order of 2 kms^{-1} , except for HD26169, for which an arc frame at that telescope position was not available and the zero point error could be as large as 15 kms^{-1} . The radial velocities which we have determined from our spectra are in good agreement with those given by Beers & Sommer-Larsen (1995) and Molaro et al. (1997), with the possible exception of HD26169, for which we note that while Beers & Sommer-Larsen (1995) give -24 kms^{-1} , Norris (1986) gives -35 kms^{-1} .

The three giants have parallaxes and proper motions measured with the Hipparcos astrometric satellite, but for the two CS stars the full kinematics cannot be recovered because there are no proper-motion data.

We used the matrices given by Johnson & Soderblom (1987) to compute spatial velocities and associated errors, adopting the left-handed reference frame in which U is directed towards the Galactic anti-center, V in the direction of Galactic rotation and W towards the North Galactic pole. We therefore changed the sign in the top row of the matrix given by Johnson & Soderblom (1987). All the velocities given are heliocentric. The error in the space velocities is dominated by the error in the parallax. The proper motion and radial velocity data yield errors in the derived space velocities of the order of a few kms^{-1} while the parallax data, yield errors which are over an order of magnitude greater. Therefore, we considered also the photometric parallax which we derived by estimating the absolute magnitude by interpolating in the proper RYI isochrone ($Y=0.25$, age of 12 Gyr and metallicity of the star) for the star's assumed effective temperature. These absolute magnitudes and the derived space velocities can be found in the Table 5.

For BD $-14^\circ 5890$ the trigonometric parallax is formally undetermined because of its large error of 126% . Its space velocities derived from the parallax listed in the Hipparcos catalogue are quite “normal” halo velocities. On the other hand, the space velocities derived from the photometric parallax are those of a runaway star, with a large retrograde orbit, which is too extreme to be believable. Even if the parallax were as large as 7 mas, i.e. 3σ larger than the Hipparcos parallax,

the kinematics of the star would still not be disc-like. Although its circular velocity would be similar to that of the sun its large U and W velocities would identify it as a halo member. We therefore consider the Hipparcos parallax to be quite close to the true parallax in spite of the large error. For HD 26169 we exclude a disc-like kinematics, for the same reasons as for BD $-14^\circ 5890$. For HD 165195 the trigonometric and photometric parallax agree within 0.03 mas. The derived space velocities differ little and qualify this star as a member of the halo. Of the three giants, however, this is the one with the smallest U and W velocities, which may suggest a disc motion. In fact if we assume the parallax to be larger by 2σ we find its rotational velocity to be 150 kms^{-1} and its kinematics may be effectively described as disc-like. HD 165195 could possess disc-like kinematics if its parallax were some 2σ larger than the Hipparcos one, however the good coincidence between the photometric and trigonometric parallaxes make this event unlikely.

We next investigated how many of the 44 stars given in Table 8 of Beers & Sommer-Larsen (1995) have Hipparcos data. We found 23 objects. We removed two from our sample because they have a negative parallax.

The kinematics of HD 111721 is quite remarkable, being in retrograde motion with a speed in excess of 600 kms^{-1} . For the three RR Lyrae variables in this list the full space kinematics have already been computed by Martin & Morrison (1998), using proper motions from Hipparcos combined with ground-based data, and distances from Layden (1994) which relied on an absolute magnitude calibration of the RR Lyrae variables. We used their data to compute the space velocities of these stars and confirm their conclusion that XX And and SW Aqr belong to the halo while EZ Lyr belongs to the metal weak thick disc. These data are also given in table 5. We draw essentially the same conclusion if we use the radial velocities from Beers & Sommer-Larsen (1995) and the Hipparcos parallaxes, except for EZ Lyr, whose Hipparcos parallax is undetermined (error larger than parallax). A caveat for RR Lyrae stars is that the radial velocity of a pulsating star is prone to systematic errors. We note here that the Simbad data base gives the values of -5 kms^{-1} and $+21 \text{ kms}^{-1}$ for SW Aqr and the values -25 kms^{-1} and -0.6 kms^{-1} for XX And.

We also reconsider the kinematics of the stars of Carney et al. (1997) since all of their “candidate thick disc” stars now have Hipparcos data. The space velocities and kinematical data are reported in Table 5. Two stars (G023-014 and G190-015) show disc-like kinematics, two (BD $+80^\circ 245$ and G182-031) have halo kinematics while one, G090-003, shows undetermined kinematics, since its Hipparcos parallax is undetermined (error larger than the parallax).

On the basis of the spectroscopically determined gravity Carney et al. computed space motions which are in substantial agreement with those given here for BD $+80^\circ 245$, G182-031 and G190-015, but in strong disagreement for G023-014.

By combining the Beers & Sommer-Larsen and the Carney et al. samples we compile a sample of candidate metal-weak thick-disc stars which possess Hipparcos proper motions using the following criteria: we exclude all stars whose Hipparcos parallax has an error greater than 200 %, but include the RR Lyrae variables using the Layden (1994) distances and the Martin & Morrison (1998) proper motions, which are based also on Hipparcos data. For these stars we adopted as errors on the space velocities $(66, 66, 52) \text{ kms}^{-1}$, which is $\sqrt{22} \times \Delta$, where 22 is the size of their metal weak disc sample (sample DISK2A) and Δ are the errors of the mean velocities of the sample quoted in their table 3. This constitutes a sample of 25 stars for which absolute space motions can be computed.

Figure 3 is a U, V diagram for all those stars and includes the three giants investigated in this paper. It can be seen that the majority of the stars qualify as true halo members. To select a sub-sample of likely metal-weak thick-disc stars we must assume a specific set of kinematic parameters: we adopt the parameters of the sample DISK2A of Martin & Morrison (1998) i.e. $(\langle U \rangle, \langle V \rangle, \langle W \rangle) = (12 - 59, -19) \text{ kms}^{-1}$ and $(\sigma(U), \sigma(V), \sigma(W)) = (64, 64, 62)$. As parameters we use V and $K^2 = (U^2 + W^2)$, which is a measure of the kinetic energy not associated with rotation. Clearly a star with high K^2 does not qualify as a disc star. To a good approximation we may write $\langle K^2 \rangle = \sigma^2(U) + \langle U \rangle^2 + \sigma^2(W) + \langle W \rangle^2$ and $\sigma^2(K^2) = 4 \langle U \rangle^2 \sigma^2(U) + 4 \langle W \rangle^2 \sigma^2(W)$. By imposing a 2σ criterion (we select all stars with $K^2 \in [2299, 12311] \text{ km}^2 \text{ s}^{-2}$ and $V \in [-187, -69] \text{ kms}^{-1}$) we isolate 7 stars: G190-15, HD165195 HD6446, HD23592, HD27928, HD97320, EZ Lyr. Only 3 stars match a 1σ criterion: G190-15, HD97320, EZ Lyr.

With such a small sample very little can be said about the nature of the metal-weak thick-disc. One of the main conclusions is that Table 8 of Beers & Sommer-Larsen (1995) is probably not the best place to look for metal-weak thick disc-stars. Quite likely the problem of finding such stars cannot be addressed with only incomplete kinematic information. In spite of this it has been possible to show that stars with disc kinematics and metallicity below $[\text{Fe}/\text{H}] \approx -1.0$ do exist. HD 166913 ($[\text{Fe}/\text{H}] = -1.80$) and HD 205650 ($[\text{Fe}/\text{H}] = -1.30$) are an other two metal-poor stars with disc kinematics which have been noted by Romano et al. (1999) in the course of their analysis of the sample of stars with Li measurements.

To demonstrate the situation we show a plot of $[\text{Fe}/\text{H}]$ versus K^2 in Fig. 4. With such a diagram, once the effects of halo contamination have been understood, and with a significant number of stars one should be able to infer knowledge of the coupling of chemical and kinematical evolution of the thick disc. A uniform distribution of stars with $[\text{Fe}/\text{H}]$ would support a slow formation of the thick disc, while a distribution peaked at some narrow range in $[\text{Fe}/\text{H}]$ would testify in favour of a short timescale.




Figure 3. U, V diagram for our sample of stars, the three giants for which abundance analysis is reported in the paper are identified by crossed symbols




Figure 4. $K^2 (= U^2 + W^2)$ as a function of $[Fe/H]$. Such a diagram should help to infer knowledge on the coupling of chemical and dynamical evolution of the thick disc.

7 CONCLUSIONS

Our analysis of chemical abundances has confirmed that all of our program stars are indeed metal-poor with iron abundances less than about one hundredth the solar abundance. The Hipparcos data allowed us to compute the kinematics of the giant stars which qualifies them as halo members; for the two fainter stars CS 22894-019 and CS 29529-012 no proper motion data are available and their membership of the thick disc cannot be established or rejected.

The ionization-equilibrium gravities are not compatible with a TO status of CS 22894-019 and CS 29529-012. On the other hand, their Li abundances prohibit their interpretation as HB stars. Therefore they are probably in the SG phase.

The chemical composition of four out of our five program stars is that typical of halo stars. CS 29529-012 is C enhanced and has a high $[Ba/Sr]$ ratio, which features make it akin to the metal-poor carbon-rich stars studied by Norris et al. (1997a). It also displays a huge Na abundance which, however, we regard as doubtful, in view of the non-detection of the subordinate Na I lines. Further observations are encouraged; it is possible that the combined effects of NLTE and a higher gravity may result in a value of $[Na/Fe]$ lower by about one dex.

In order to shed further light on the metal-weak thick-disc, proper motion data for these two stars, as well as for the other




Figure 5. The Ba resonance line in CS 29529-12, the dotted line is a synthetic spectrum computed assuming $\epsilon = +0.48$.

metal-poor stars of comparable magnitude are badly needed. The present paper, in line with other recent chemical analyses of thick disc stars (Nissen & Schuster 1997, Fuhrmann 1998a), suggests that no chemical signature is likely to be found to discriminate between halo and metal-weak thick-disc.

ACKNOWLEDGMENTS

We are grateful to J. Danziger for critically reading our manuscript and suggesting many improvements. Special thanks are due to T. Valente for his help in the measuring of equivalent widths. Use was made of the SIMBAD data base, operated at the CDS, Strasbourg, France. This research was partially supported by Collaborative NATO grant No. 950875. This research is based on observations collected at the European Southern Observatory, Chile.

REFERENCES

- Alonso A., Arribas S., Martínez-Roger C., 1996b, *A&A* in press, IAC preprint
Baumüller D., Butler K., Gehren T., 1998, *A&A* 338, 661
Beers T. B., Sommer-Larson J., 1995, *ApJS* 96, 175
Beers T.C., Preston G.W., Shectman S.A., 1992, *AJ* 103, 1987
Bertelli G., Bressan A., Chiosi C., Fagotto F., Nasi E., 1994, *A&AS* 106, 275
Bonifacio P., Molaro P., Beers T.C., Vladilo G., 1998, *A&A* 332, 672
Carney B.W., Latham D.W., Laird J.B., 1989, *AJ* 97, 423
Carney B.W., Laird J.B., Latham D.W., Aguilar L.A., 1996, *AJ* 112, 668
Carney B.W., Wright J.S., Sneden C., Laird J.B., Aguilar L.A., Latham D.W., 1997, *AJ* 114, 363
Cavallo R.M., Pilachowski C.A., Rebolo R., 1997, *PASP* 109, 226
Cayrel R., 1988 in Cayrel de Strobel G., Spite M., eds, *Proc. IAU Symp. 132, The Impact of Very High S/N Spectroscopy on Stellar Physics*. Kluwer, Dordrecht, p. 345
Francois P., 1996, *A&A* 313, 229
Friel E.D., 1987, *AJ* 93, 1388
Fuhrmann K., 1998a, *A&A* 338, 161
Fuhrmann K., 1998b, *A&A* 330, 626
Gilmore G., Reid N., 1983, *MNRAS* 202, 1025
Gilmore G., Wyse R.F.G., 1985, *AJ* 90, 2015
Gilmore G., Wyse R.F.G., Kuijken K., 1989, *ARA&A* 27, 555
Gilroy K.W., Sneden C., Cowan J.J., 1988, *ApJ* 327, 298
Gratton R., Sneden C., 1991, *A&A* 241, 501
Green E.M., Demarque P., King C.R., 1987 *The Revised Yale Isochrones and Luminosity Functions* (Yale Observatory, New Haven), made available electronically by the Centre D'observatoire Stellaires, Strasbourg
Hartkopf W.L., Yoss K.M., 1982, *AJ* 87, 1679
Jehin E., Magain P., Neuforge C., Noels A., Thoul A.A., 1998, *A&A* 330, L33
Johnson D.R.H., Soderblom D.R., 1987, *AJ* 93, 864

Table 2.(continued) line data and individual abundances (continued)

	$\lambda(\text{nm})$	$\chi(\text{cm}^{-1})$	loggf	$EW(pm)$ CS 22894-19	ϵ	$EW(pm)$ CS 29529-12	ϵ	$EW(pm)$ BD -14°5890	ϵ	$EW(pm)$ HD 165195	ϵ	$EW(pm)$ HD 26169	ϵ
Fe I	400.5241	12560.930	-0.570	7.10	4.72	—	—	—	—	—	—	10.88	4.69
Fe I	404.5813	11976.234	0.220	10.17	4.58	11.45	5.35	—	—	—	—	17.46	4.92
Fe I	406.3594	12560.930	0.080	8.48	4.39	8.92	4.86	18.18	4.92	—	—	14.76	4.81
Fe I	407.1737	12968.549	0.000	8.25	4.45	—	—	18.69	5.09	—	—	14.44	4.90
Fe I	413.2058	12968.549	-0.634	—	—	—	—	13.42	5.01	19.02	5.59	11.24	4.85
Fe I	414.3869	12560.930	-0.440	6.58	4.47	7.71	5.01	—	—	—	—	—	—
Fe I	418.7795	43434.629	-0.520	—	—	—	—	10.71	5.22	12.85	5.69	8.53	4.95
Fe I	419.8304	43163.327	-0.521	2.32	4.52	5.52	5.35	—	—	—	—	9.25	5.09
Fe I	419.9095	24574.650	0.248	3.51	4.61	—	—	9.96	5.02	—	—	7.60	4.65
Fe I	420.2028	11976.234	-0.697	5.94	4.53	—	—	12.77	4.76	18.40	5.47	—	—
Fe I	423.3602	43633.534	-0.560	2.67	4.72	3.68	5.08	9.73	5.06	13.39	5.86	8.50	5.02
Fe I	423.5936	43163.327	-0.292	—	—	7.52	5.69	—	—	—	—	—	—
Fe I	425.0118	43434.629	-0.370	2.69	4.52	5.40	5.23	9.23	4.73	12.04	5.38	7.90	4.67
Fe I	425.0787	12560.930	-0.691	—	—	6.70	4.99	12.55	4.78	17.50	5.45	11.10	4.78
Fe I	426.0473	42815.858	0.129	6.22	4.61	8.26	5.36	—	—	—	—	—	—
Fe I	427.1153	19757.033	-0.334	3.63	4.65	6.89	5.50	10.26	4.92	11.41	5.17	9.11	4.91
Fe I	427.1759	11976.234	-0.127	8.19	4.42	8.87	4.93	—	—	21.38	5.07	—	—
Fe I	428.2402	17550.175	-0.730	2.77	4.62	5.45	5.33	10.93	5.20	—	—	9.25	5.04
Fe I	430.7902	12560.930	-0.060	9.80	4.79	—	—	—	—	—	—	—	—
Fe I	432.5762	12968.549	0.000	8.54	4.47	10.73	5.43	19.66	5.13	—	—	14.60	4.85
Fe I	437.5930	0.000	-3.031	1.74	4.58	—	—	10.81	4.75	16.23	5.86	9.55	4.91
Fe I	438.3544	12560.930	0.160	10.37	4.62	11.72	5.48	19.68	4.79	—	—	16.18	4.76
Fe I	440.4750	12560.930	-0.180	9.49	4.82	8.43	4.91	17.68	5.03	—	—	14.33	4.89
Fe I	441.5122	12968.549	-0.580	7.40	4.79	7.44	5.09	—	—	—	—	11.62	4.79
Fe I	452.8613	17550.175	-0.822	3.87	4.91	3.92	5.10	—	—	—	—	9.83	5.16
Fe I	489.1492	43434.629	-0.130	3.22	4.71	5.12	5.24	10.59	5.09	13.35	5.61	8.84	4.97
Fe I	492.0502	43163.327	0.070	3.30	4.51	6.94	5.41	11.25	5.01	16.80	5.89	9.87	4.97
Fe I	495.7298	43163.327	-0.342	—	—	4.97	5.42	9.71	5.09	18.14	5.90	7.54	4.90
Fe I	495.7597	42815.858	0.127	—	—	—	—	—	—	—	—	10.77	5.09
Fe I	522.7189	12560.930	-0.969	4.24	4.49	—	—	14.43	5.14	19.24	5.73	11.86	5.01
Fe I	523.2939	42815.858	-0.140	3.34	4.81	5.67	5.42	10.41	5.10	13.15	5.59	8.42	4.96
Fe II	423.3172	20830.582	-2.000	2.48	4.64	4.39	4.92	9.04	5.02	—	—	7.65	4.94
Fe II	455.5893	22810.357	-2.290	—	—	3.58	5.25	—	—	—	—	—	—
Fe II	492.3927	23317.633	-1.320	4.32	4.57	8.02	5.22	10.84	4.95	12.80	5.57	9.19	4.83
Fe II	501.8440	23317.633	-1.220	5.12	4.60	9.81	5.53	12.45	5.14	14.83	5.72	10.03	4.89
Fe II	516.9033	23317.633	-0.870	—	—	9.81	5.17	—	—	—	—	—	—
Sr II	407.7709	0.000	0.167	6.90	-0.40	6.81	-0.33	17.68	0.25	20.37	0.50	17.08	0.55
Sr II	421.5519	0.000	-0.145	6.00	-0.29	6.90	-0.02	14.71	0.12	19.47	0.70	14.74	0.51
Ba II	455.4029	0.000	0.170	< 1.3	< -1.4	9.34	0.48	13.16	-0.34	11.08	-0.67	11.08	-0.34

Table 3. Errors on abundances for CS29529-12

Element	σ	$\Delta_{\log g+0.5}$	Δ_{T-130K}	$\Delta_{\xi+0.3\text{km/s}}$	$\Delta_{EW+0.4\text{pm}}$
Na	0.08	-0.31	-0.26	-0.25	+0.09
Mg	0.17	-0.15	-0.11	-0.08	+0.05
Al	—	-0.01	-0.11	-0.15	+0.09
Ca	0.15	-0.11	-0.17	-0.09	+0.08
Sc	—	+0.17	-0.06	-0.10	+0.09
Ti	0.20	+0.17	-0.05	-0.12	+0.10
Cr	0.16	-0.01	-0.12	-0.06	+0.08
Fe	0.21	+0.02	-0.11	-0.13	+0.09
Sr	—	+0.15	-0.09	-0.14	+0.12
Ba	—	+0.12	-0.08	-0.27	+0.11

Kurucz R.L., 1993, CD-ROM No. 13, 18

Layden A.C., 1994, AJ 108, 1016

Luck R.E., Bond H.E., 1982, ApJ 259, 792

Martin J.C., Morrison H.L., 1998, AJ 116, 1724

Majewski S.R., 1992, ApJS 78, 87

McWilliam A., Preston G.W., Sneden C., Searle L., 1995, AJ 109, 2757

Table 4. Abundances

	$\epsilon_{\odot}$	ϵ	[X/Fe]	ϵ	[X/Fe]	ϵ	[X/Fe]	ϵ	[X/Fe]	ϵ	[X/Fe]
	CS 22894-19			CS 29529-12		BD -14° 5890		HD 165195		HD 26169	
Na	6.33	3.13	-0.29	5.62 :	1.56 :						
Mg	7.58	5.04	$+0.36$	5.85	$+0.54$	5.61	$+0.55$	5.74	$+0.08$	5.46	$+0.49$
Al	6.47	3.01	-0.56	3.70	-0.50	3.14	-0.81	3.90	-0.65	3.16	-0.72
Ca	6.36	3.80	$+0.34$	4.41	$+0.32$	4.10	$+0.26$	—	—	4.07	$+0.32$
Sc	3.10	0.07	-0.13	0.74	-0.09	0.42	-0.16	1.12	-0.06	0.44	-0.07
Ti	4.99	2.36	$+0.27$	3.03	$+0.31$	2.70	$+0.23$	3.60	$+0.53$	2.63	$+0.25$
Cr	5.67	2.64	-0.13	3.43	$+0.03$	2.90	-0.25	3.50	-0.25	2.72	-0.36
Mn	5.39	2.18	-0.31	—	—	3.06	$+0.19$	—	—	2.68	-0.12
Fe	7.51	4.61	0.00	5.24	0.00	4.99	0.00	5.59	0.00	4.90	0.00
Sr	2.90	-0.34	-0.34	-0.02	-0.65	0.18	-0.20	0.61	-0.37	0.53	$+0.24$
Ba	2.13	< -1.40	< -0.60	$+0.48$	$+0.62$	-0.34	$+0.05$	-0.67	-0.88	-0.34	$+0.13$

a : denotes an uncertain value

Table 5. kinematical data

Star	π mas	μ_{α} mas	μ_{δ} mas	v_r kms $^{-1}$	U kms $^{-1}$	V kms $^{-1}$	W kms $^{-1}$
BD -14° 5890	1.48 ± 1.86 0.57^a	-16.96 ± 2.05	-83.97 ± 1.67	+118	-205 ± 158 -406	-183 ± 300 -564	-116 ± 63 -196
HD 165195	2.20 ± 1.04 2.34^a	-28.55 ± 0.83	-82.87 ± 0.68	0	-102 ± 48 -96	-157 ± 74 -148	-26 ± 13 -26
HD 26169	2.83 ± 0.79 1.95^a	137.18 ± 0.80	74.90 ± 1.04	-20	203 ± 55 292	-129 ± 42 -194	106 ± 28 148
HD6446	1.96 ± 0.57	38.98 ± 0.53	-48.69 ± 0.59	62	6 ± 10	-153 ± 33	55 ± 28
XX And ^b	1.50 ± 2.15 1.07	58.97 ± 1.64 55.76	-35.46 ± 1.47 -35.16	-25 44	118 ± 189 118	-170 ± 218 -170	-72 ± 117 -72
HD11569	9.47 ± 0.77	268.97 ± 0.71	-29.45 ± 0.93	-18	97 ± 8	-80 ± 10	54 ± 8
HD23592	1.67 ± 1.00	14.37 ± 0.98	25.68 ± 1.06	-54	78 ± 50	36 ± 7	50 ± 9
HD27928	2.36 ± 1.16	26.90 ± 0.90	-67.47 ± 1.22	15	-98 ± 51	-107 ± 48	22 ± 17
HD33771	1.64 ± 1.08	79.65 ± 0.99	-30.25 ± 1.11	18	-45 ± 35	-179 ± 110	164 ± 115
CD-33 3337	9.11 ± 1.01	-178.43 ± 0.90	-148.74 ± 0.97	62	-15 ± 6	-44 ± 9	-128 ± 13
HD81223	3.39 ± 0.69	-102.14 ± 0.82	85.68 ± 0.66	-52	193 ± 37	4 ± 13	-7 ± 5
HD83212	1.96 ± 0.98	-15.51 ± 0.92	-20.41 ± 0.65	110	26 ± 3	-123 ± 16	-14 ± 29
HD97320	17.77 ± 0.76	159.19 ± 0.74	-201.28 ± 0.74	48	-72 ± 5	-18 ± 9	-38 ± 2
BD-01 2582	2.98 ± 1.35	-21.19 ± 1.34	-117.77 ± 0.78	0	-61 ± 28	-151 ± 69	-99 ± 46
HD111721	3.29 ± 1.11	-273.70 ± 0.94	-321.99 ± 0.66	24	131 ± 47	-524 ± 173	-282 ± 101
HD118055	$.66 \pm 1.24$	-23.09 ± 1.64	-14.69 ± 0.94	-101	145 ± 172	-121 ± 317	-115 ± 82
HD122196	9.77 ± 1.32	-452.86 ± 1.08	-82.55 ± 1.01	-24	172 ± 22	-144 ± 22	16 ± 5
HD136316	1.08 ± 0.95	-27.55 ± 0.86	-64.44 ± 0.83	-45	178 ± 125	-189 ± 189	-171 ± 149
EZ Lyr ^b	0.95	-1.59	13.10	-60.00	95	-33	24
SW Aqr ^b	1.15 ± 2.46 0.74	-42.05 ± 2.63 -42.89	-58.87 ± 1.96 -59.11	-62 -36	-207 ± 511 -274	-219 ± 381 -240	44 ± 27 36
G090-003	1.20 ± 1.41	16.93 ± 2.36	-199.64 ± 2.44	29	14 ± 26	-774 ± 1516	-168 ± 352
BD+80245	3.91 ± 0.88	136.81 ± 1.22	-366.66 ± 0.99	4	183 ± 56	-366 ± 115	241 ± 75
G182-031	5.91 ± 0.98	-154.86 ± 1.17	-244.96 ± 1.16	-61	-158 ± 36	-178 ± 27	32 ± 13
G023-014	8.11 ± 4.40	-140.42 ± 6.21	-231.74 ± 7.71	18	-126 ± 87	-98 ± 85	1 ± 6
G190-015	14.09 ± 1.27	175.38 ± 2.20	-313.55 ± 1.86	-55	-8 ± 2	-93 ± 11	-95 ± 18

^a photometric, estimated from M_v derived from isochrones^b distance from Layden (1994), proper motions from Martin&Morrison (1998)

Molaro, P., Bonifacio P, Castelli F., Pasquini L.,1997, A&A 319, 593

Molaro P., Bonifacio P., Pasquini L.,1997, MNRAS 292, L1

Morrison H.L., Flynn C., Freeman K.C.,1990, AJ 100, 1191

Nissen P.E., Schuster W.J.,1997, A&A 326, 751

Nissen P.E., Hoeg E., Schuster W.J., 1997 in Proceedings of the ESA Symposium “Hipparcos - Venice ’97”, SP-402, p. 225

Norris J.E.,1986, ApJS 61, 667

Norris J. E., Ryan S.G.,1991, ApJ 380, 403

Norris J.E., Bessel M.S., Pickles A.J.,1985, ApJS 58, 463

- Norris J.E., Ryan S.G., Beers T.C., 1997a, ApJ 488, 350
 Norris J.E., Ryan S.G., Beers T.C., 1997b, ApJ 489, L169
 Peterson R.C., Kurucz R.L., Carney B.W., 1990, ApJ 350, 173
 Pilachowski C.A., Sneden C., Booth J., 1993, ApJ 407, 699
 Preston G. W., Beers T., C., Shectman S. A., 1994, AJ 108, 538
 Primas F., Molaro P., Castelli F., 1994, A&A 290, 885
 Ratnatunga K.U., Freeman K.C., 1989, ApJ 339, 126
 Reid I.N., 1998, AJ 115, 204
 Reid I.N., Gizis, J.E., Coehen J.G., Pahre M.A., Hogg D.W., Cowie L., Hu E., Songaila A., 1997, PASP 109, 559
 Romano D., Matteucci F., Molaro P., Bonifacio P., 1999, A&A , in press
 Ryan S.G., Lambert D.L., 1995, AJ 109, 2068
 Ryan S.G., Norris J.E., Beers T.C., 1996, ApJ 471, 254
 Sandage A., Fouts G., 1987, AJ 93, 74
 Timmes F.X., Woosley S.E., Weaver T.A., 1995, ApJS 98, 617
 Van der Kruit P.C., Searle L., A&A 1981, 95105
 Van der Kruit P.C., Searle L., A&A 1981, 95116
 Van der Kruit P.C., Searle L., A&A 1981, 11061
 Wyse R.F.G., Gilmore G., 1995, AJ 110, 2771
 Yoss K.M., Neese C.L., Hartkopf, W.I., 1987, AJ 94, 1600
 Younger S.M., Fuhr J.R., Martin G.A., Wiese W.L., 1978 J. of Phys. and Chem. Ref. Data, 7 495

APPENDIX A: COMPARISON WITH PREVIOUS RESULTS

For BD-14°5890, the only high-resolution studies which allow a comparison to are those of Pilachowski, Sneden & Booth (1993) and Cavallo, Pilachowski & Rebolo (1997). The [Fe/H] and [Ca/H] of Pilachowski et al. (1993) are 0.45 dex and 0.54 dex respectively higher than ours. This discrepancy can be largely attributed to the difference in the adopted effective temperatures: our T_{eff} is 250 K cooler. The [Fe/H] of Cavallo et al. is 0.26 dex higher than ours and in this case the main cause of the discrepancy is the different microturbulent velocity. However, if we adopt the value of Cavallo et al. (0.8 km s^{-1}) our [Fe/H] becomes 0.4 dex *larger* than that of Cavallo et al. ; incidentally our line to line scatter doubles with this low microturbulent velocity.

For HD 165195 we compared our results with those of Francois (1996), Gratton & Sneden (1991) and Gilroy et al. (1988). With respect to Francois our [Fe/H] is 0.25 dex larger, and this may be attributed to our lower microturbulent velocity. In contrast there is a strong disagreement in the Ba abundance, ours is about 0.9 dex *lower* than that of Francois. We have no explanation for this difference, but point out that the result of Francois was based only on the 614.17 nm line while ours only on the 455.4 nm line. With Gratton & Sneden the agreement is generally good at the 0.1 - 0.2 dex level, with the exception of Ti where our abundance is over 0.3 dex larger. Also in this case our lower microturbulent velocity is the main responsible for the discrepancy. Also the comparison with Gilroy et al. is generally good, again with the notable exception of Ti (ours is 0.66 dex larger), The microturbulence adopted by Gilroy et al. is the highest for this star among the reviewed studies: 2.8 km s^{-1} . With this high value of ξ our data imply a Ti abundance which is 0.73 dex lower, i.e. in substantial agreement with that of Gilroy et al. .

For HD 26169 we compared our results with those of Peterson, Kurucz & Carney (1990); for all the elements in common our abundances are lower than those of Peterson et al. , only a few hundredths of dex for Sc and Ba and as much as 0.36 dex for Cr. While this agreement may be regarded as satisfactory in the context of the claimed accuracy of the present analysis, it is somewhat disappointing, considering the fact that we have virtually the same atmospheric parameters (the only difference is 0.1 km s^{-1} in ξ) and adopt the same kind of models.

This paper has been produced using the Royal Astronomical Society/Blackwell Science $\text{\TeX}$ macros.